Test Booklet Code

KANHA

No.:

F1

This Booklet contains 24 pages.

Do not open this Test Booklet until you are asked to do so.

Important Instructions:

- 1. The Answer Sheet is inside this Test Booklet. When you are directed to open the Test Booklet, take out the Answer Sheet and fill in the particulars on **side-1** and **side-2** carefully with **blue/black** ball point pen only.
- 2. The test is of **3 hours** duration and Test Booklet contains **180** questions. Each question carries **4** marks. For each correct response, the candidate will get **4** marks. For each incorrect response, **one mark** will be deducted from the total scores. The maximum marks are **720**.
- 3. Use Blue/Black Ball Point Pen only for writing particulars on this page/marking responses.
- 4. Rough work is to be done on the space provided for this purpose in the Test Booklet only.
- 5. On completion of the test, the candidate must hand over the Answer Sheet to the invigilator before leaving the Room/Hall. The candidates are allowed to take away this Test Booklet with them.
- 6. The CODE for this Booklet is **F1**. Make sure that the CODE printed on **Side-2** of the Answer Sheet is the same as that on this Test Booklet. In case of discrepancy, the candidate should immediately report the matter to the Invigilator for replacement of both the Test Booklet and the Answer Sheet.
- 7. The candidates should ensure that the Answer Sheet is not folded. Do not make any stray marks on the Answer Sheet. Do not write your Roll No. anywhere else except in the specified space in the Test Booklet/Answer Sheet.
- 8. Use of white fluid for correction is **NOT** permissible on the Answer Sheet.
- 9. Each candidate must show on demand his/her Admit Card to the Invigilator.
- 10. No candidate, without special permission of the Superintendent or Invigilator, would leave his/her seat.
- 11. The candidates should not leave the Examination Hall without handing over their Answer Sheet to the Invigilator on duty and sign the Attendance Sheet twice. Cases where a candidate has not signed the Attendance Sheet second time will be deemed not to have handed over the Answer Sheet and dealt with as an unfair means case.
- 12. Use of Electronic/Manual Calculator is prohibited.
- 13. The candidates are governed by all Rules and Regulations of the examination with regard to their conduct in the Examination Hall. All cases of unfair means will be dealt with as per Rules and Regulations of this examination.
- 14. No part of the Test Booklet and Answer Sheet shall be detached under any circumstances.
- 15. The candidates will write the Correct Test Booklet Code as given in the Test Booklet/Answer Sheet in the Attendance Sheet.

Name of the Car	ndidate (in Capitals) :	
Roll Number	: in figures	
Ivoir I valificor	: in words	
Centre of Exami		
	, , ,	Invigilator's Signature :
Facsimile signat		
_	endent:	

- 1. In light reaction, plastoquinone facilitates the | 6. transfer of electrons from:
 - PS-I to NADP+ (1)
 - (2)PS-I to ATP synthase
 - PS-II to Cytb₆f complex (3)
 - Cytb₆f complex to PS-I (4)
- 2. The sequence that controls the copy number of the linked DNA in the vector, is termed:
 - (1) Palindromic sequence
 - (2)Recognition site
 - (3)Selectable marker
 - (4) Ori site
- The specific palindromic sequence which is 3. recognized by EcoRI is:
 - 5' CTTAAG 3' (1)
 - 3' GAATTC 5'
 - 5' GGATCC 3' (2)
 - 3' CCTAGG 5'
 - 5' GAATTC 3' (3)
 - 3' CTTAAG 5'
 - **(4)** 5' - GGAACC - 3'
 - 3' CCTTGG 5'
- 4. Identify the **wrong** statement with reference to immunity.
 - (1) Active immunity is quick and gives full response.
 - Foetus receives some antibodies from (2)mother, it is an example for passive immunity.
 - (3)When exposed to antigen (living or dead) antibodies are produced in the host's body. It is called "Active immunity".
 - When ready-made antibodies are directly (4)given, it is called "Passive immunity".
- Experimental verification of the chromosomal **5**. theory of inheritance was done by:
 - Boveri (1)
 - (2)Morgan
 - (3)Mendel
 - Sutton (4)

- Match the following concerning essential elements and their functions in plants:
 - Iron
- (i) Photolysis of water
- (b) Zinc
- Pollen germination (ii)
- (c) Boron
- Required for chlorophyll (iii) biosynthesis
- Manganese (iv) IAA biosynthesis

Select the **correct** option:

- (a) (b)
- (iii) (iv)
- (ii)

(d)

(i)

(iii)

(iii)

(c)

(2)(iv) (i)

(iv)

(1)

(3)

(4)

- (ii) (iv)
- (ii) (i)
 - (iii)
- (i) (ii)
- 7. In gel electrophoresis, separated DNA fragments can be visualized with the help of:
 - (1) Acetocarmine in UV radiation
 - Ethidium bromide in infrared radiation (2)
 - (3)Acetocarmine in bright blue light
 - (4) Ethidium bromide in UV radiation
- 8. Name the enzyme that facilitates opening of DNA helix during transcription.
 - (1) DNA polymerase
 - (2)RNA polymerase
 - **DNA** ligase (3)
 - (4) DNA helicase
- 9. In which of the following techniques, the embryos are transferred to assist those females who cannot conceive?
 - ICSI and ZIFT (1)
 - (2)GIFT and ICSI
 - ZIFT and IUT (3)
 - GIFT and ZIFT (4)
- 10. Identify the basic amino acid from the following.
 - Lysine (1)
 - (2)Valine
 - (3)Tyrosine
 - (4) Glutamic Acid

- 11. Identify the **wrong** statement with reference to transport of oxygen.
 - (1) Higher H^+ conc. in alveoli favours the formation of oxyhaemoglobin.
 - (2) Low pCO_2 in alveoli favours the formation of oxyhaemoglobin.
 - (3) Binding of oxygen with haemoglobin is mainly related to partial pressure of O_2 .
 - (4) Partial pressure of CO_2 can interfere with O_2 binding with haemoglobin.
- 12. Floridean starch has structure similar to:
 - (1) Mannitol and algin
 - (2) Laminarin and cellulose
 - (3) Starch and cellulose
 - (4) Amylopectin and glycogen
- **13.** By which method was a new breed 'Hisardale' of sheep formed by using Bikaneri ewes and Marino rams?
 - (1) Cross breeding
 - (2) Inbreeding
 - (3) Out crossing
 - (4) Mutational breeding
- 14. Match the following columns and select the **correct** option.

Column - II Column - I (a) Pituitary gland (i) Grave's disease Thyroid gland (b) (ii) Diabetes mellitus Adrenal gland (c) (iii) Diabetes insipidus (d) Pancreas (iv) Addison's disease (a) (b) **(c)** (d) (1) (iii) (i) (iv) (ii) (2)(ii) (i) (iv) (iii) (3)(iv) (iii) (i) (ii) (4) (iii) (ii)(i) (iv)

- **15.** Select the option including all sexually transmitted diseases.
 - (1) AIDS, Malaria, Filaria
 - (2) Cancer, AIDS, Syphilis
 - (3) Gonorrhoea, Syphilis, Genital herpes
 - (4) Gonorrhoea, Malaria, Genital herpes

- **16.** Choose the **correct** pair from the following:
 - $\begin{array}{ccc} \hbox{(1)} & \hbox{Nucleases} & \hbox{-} & \hbox{Separate the two strands} \\ & \hbox{of DNA} \end{array}$
 - (2) Exonucleases Make cuts at specific positions within DNA
 - (3) Ligases Join the two DNA molecules
 - $\begin{array}{ccc} \text{(4)} & \text{Polymerases -} & \text{Break the DNA into} \\ & & \text{fragments} \end{array}$
- **17.** Ray florets have:
 - (1) Hypogynous ovary
 - (2) Half inferior ovary
 - (3) Inferior ovary
 - (4) Superior ovary
- 18. Match the organism with its use in biotechnology.
 - (a) Bacillus (i) Cloning vector thuringiensis
 - (b) Thermus (ii) Construction of aquaticus first rDNA molecule
 - $\begin{array}{ccc} \mbox{(c)} & A grobacterium & \mbox{(iii)} & \mbox{DNA polymerase} \\ & tume faciens \end{array}$
 - $\begin{array}{ccc} \text{(d)} & Salmonella & \text{(iv)} & \text{Cry proteins} \\ & & typhimurium \end{array}$

Select the **correct** option from the following:

- (a) (b) **(c)** (d) (iii) (1) (ii) (iv) (i) (2)(iii) (iv) (i) (ii) (3)(ii) (iv) (iii) (i) (4) (iv) (iii) (i) (ii)
- **19.** The product(s) of reaction catalyzed by nitrogenase in root nodules of leguminous plants is/are:
 - (1) Ammonia and oxygen
 - (2) Ammonia and hydrogen
 - (3) Ammonia alone
 - (4) Nitrate alone

spra	ne the plant growth regulator which upon aying on sugarcane crop, increases the length tem, thus increasing the yield of sugarcane o.
(1)	Ethylene
(2)	Abscisic acid
(3)	Cytokinin
(4)	Gibberellin
The	body of the ovule is fused within the funicle
(1)	Nucellus
(2)	Chalaza
(3)	Hilum
(4)	Micropyle
The	process of growth is maximum during:
(1)	Senescence
(2)	Dormancy
(3)	Logphase
(4)	Lag phase
	aterally symmetrical and acoelomate animals exemplified by :
(1)	Aschelminthes
(2)	Annelida
(3)	Ctenophora
(4)	Platyhelminthes
	ich of the following is put into Anaerobic sludge ester for further sewage treatment?
(1)	Effluents of primary treatment
(2)	Activated sludge
(3)	Primary sludge
(4)	Floating debris

25. Match the following columns and select the **correct** option.

	Colı	ımn -	I		Column - II
(a)	Floa	ting Ri	bs	(i)	Located between
					second and
					seventh ribs
(b)	Acro	mion		(ii)	Head of the
					Humerus
(c)	Scap	ula		(iii)	Clavicle
(d)	Glen	oid cav	vity	(iv)	Do not connect
					with the sternur
	(a)	(b)	(c)	(d)	
(1)	(iii)	(ii)	(iv)	(i)	
(2)	(iv)	(iii)	(i)	(ii)	
(3)	(ii)	(iv)	(i)	(iii)	
(4)	(i)	(iii)	(ii)	(iv)	

- **26.** Identify the **wrong** statement with regard to Restriction Enzymes.
 - (1) They are useful in genetic engineering.
 - (2) Sticky ends can be joined by using DNA ligases.
 - (3) Each restriction enzyme functions by inspecting the length of a DNA sequence.
 - (4) They cut the strand of DNA at palindromic sites.
- 27. Match the following columns and select the correct option.

	Column - I			Co	lumn - II	
(a)	Greg	arious	, polyp	hagou	s (i)	Asterias
	pest					
(b)	Adul	t with	radial		(ii)	Scorpion
	symi	metry	and la	rva		
	with	bilate	ral syn	nmetry	7	
(c)	Book	lungs			(iii)	Ctenoplana
(d)	Biolu	umines	cence		(iv)	Locusta
	(a)	(b)	(c)	(d)		
(1)	(iii)	(ii)	(i)	(iv)		
(2)	(ii)	(i)	(iii)	(iv)		
(3)	(i)	(iii)	(ii)	(iv)		
(4)	(iv)	(i)	(ii)	(iii)		

- **28.** If the head of cockroach is removed, it may live for few days because :
 - the head holds a small proportion of a nervous system while the rest is situated along the ventral part of its body.
 - (2) the head holds a $1/3^{rd}$ of a nervous system while the rest is situated along the dorsal part of its body.
 - (3) the supra-oesophageal ganglia of the cockroach are situated in ventral part of abdomen.
 - (4) the cockroach does not have nervous system.

- **29.** Which of the following regions of the globe exhibits highest species diversity?
 - (1) Himalayas
 - (2) Amazon forests
 - (3) Western Ghats of India
 - (4) Madagascar
- **30.** Which is the important site of formation of glycoproteins and glycolipids in eukaryotic cells?
 - (1) Golgi bodies
 - (2) Polysomes
 - (3) Endoplasmic reticulum
 - (4) Peroxisomes
- **31.** Which of the following pairs is of unicellular algae?
 - (1) Anabaena and Volvox
 - (2) Chlorella and Spirulina
 - (3) Laminaria and Sargassum
 - (4) Gelidium and Gracilaria
- **32.** Which one of the following is the most abundant protein in the animals?
 - (1) Lectin
 - (2) Insulin
 - (3) Haemoglobin
 - (4) Collagen
- **33.** Dissolution of the synaptonemal complex occurs during:
 - (1) Diplotene
 - (2) Leptotene
 - (3) Pachytene
 - (4) Zygotene
- **34.** How many true breeding pea plant varieties did Mendel select as pairs, which were similar except in one character with contrasting traits?
 - (1) 14
 - (2) 8
 - (3) 4
 - (4) 2

- **35.** Cuboidal epithelium with brush border of microvilli is found in :
 - (1) proximal convoluted tubule of nephron
 - (2) eustachian tube

5

- (3) lining of intestine
- (4) ducts of salivary glands
- **36.** Match the following with respect to meiosis:
 - (a) Zygotene (i) Terminalization
 - (b) Pachytene (ii) Chiasmata
 - (c) Diplotene (iii) Crossing over
 - (d) Diakinesis (iv) Synapsis

Select the **correct** option from the following:

- (a) (b) (c) (d)
- (1) (i) (ii) (iv) (iii)
- (2) (ii) (iv) (iii) (i)
- (3) (iii) (iv) (i) (ii)
- (4) (iv) (iii) (ii) (i)
- **37.** Which of the following statements about inclusion bodies is **incorrect**?
 - (1) They lie free in the cytoplasm.
 - (2) These represent reserve material in cytoplasm.
 - (3) They are not bound by any membrane.
 - (4) These are involved in ingestion of food particles.
- **38.** Which of the following would help in prevention of diuresis?
 - (1) Atrial natriuretic factor causes vasoconstriction
 - (2) Decrease in secretion of renin by JG cells
 - (3) More water reabsorption due to undersecretion of ADH
 - (4) Reabsorption of Na⁺ and water from renal tubules due to aldosterone

(d)

(1)

(2)

(3)

Malaria

(a)

(ii)

(iv)

(i)

(iii)

(b)

(i)

(i)

(iii)

(iv)

(iv)

(d)

(iv)

(iii)

(iv)

(ii)

(c)

(iii)

(ii)

(ii)

(i)

Haemophilus

(4)

(1)

(2)

(3)

(4)

42.

(iii)

(ii)

Industrial melanism

Natural selection

Adaptive radiation

Convergent evolution

(i)

(iv)

Flippers of Penguins and Dolphins are examples

- **48.** Identify the **wrong** statement with reference to the gene 'I' that controls ABO blood groups.
 - (1) When I^A and I^B are present together, they express same type of sugar.
 - (2) Allele 'i' does not produce any sugar.
 - (3) The gene (I) has three alleles.
 - (4) A person will have only two of the three alleles.
- **49.** According to Robert May, the global species diversity is about:
 - (1) 50 million
 - (2) 7 million
 - (3) 1.5 million
 - (4) 20 million
- **50.** Which of the following is **not** an attribute of a population?
 - (1) Mortality
 - (2) Species interaction
 - (3) Sex ratio
 - (4) Natality
- **51.** In water hyacinth and water lily, pollination takes place by:
 - (1) wind and water
 - (2) insects and water
 - (3) insects or wind
 - (4) water currents only
- **52.** The QRS complex in a standard ECG represents:
 - (1) Depolarisation of ventricles
 - (2) Repolarisation of ventricles
 - (3) Repolarisation of auricles
 - (4) Depolarisation of auricles
- **53.** Select the **correct** match.
 - (1) Sickle cell anaemia Autosomal recessive trait, chromosome-11
 - (2) Thalassemia X linked
 - (3) Haemophilia Ylinked
 - (4) Phenylketonuria Autosomal dominant trait

- **54.** The number of substrate level phosphorylations in one turn of citric acid cycle is :
 - (1) Two

7

- (2) Three
- (3) Zero
- (4) One
- **55.** Match the following:
 - (a) Inhibitor of catalytic (i) Ricin activity
 - (b) Possess peptide bonds (ii) Malonate
 - (c) Cell wall material in (iii) Chitin fungi
 - (d) Secondary metabolite (iv) Collagen

Choose the **correct** option from the following:

- (a) (b) (c) (d)
- (1) (iii) (iv) (i) (ii)
- (2) (ii) (iii) (i) (iv)
- (3) (ii) (iv) (iii) (i)
- (4) (iii) (i) (iv) (ii)
- **56.** Which of the following refer to **correct** example(s) of organisms which have evolved due to changes in environment brought about by anthropogenic action?
 - (a) Darwin's Finches of Galapagos islands.
 - (b) Herbicide resistant weeds.
 - (c) Drug resistant eukaryotes.
 - (d) Man-created breeds of domesticated animals like dogs.
 - (1) (b), (c) and (d)
 - (2) only (d)
 - (3) only (a)
 - (4) (a) and (c)
- 57. Some dividing cells exit the cell cycle and enter vegetative inactive stage. This is called quiescent stage (G_0). This process occurs at the end of:
 - (1) Sphase
 - G_2 phase
 - (3) M phase
 - (4) G_1 phase

68 .	The roots that originate from the base of the stem
	are:

- (1) Prop roots
- (2) Lateral roots
- (3) Fibrous roots
- (4) Primary roots
- **69.** Goblet cells of alimentary canal are modified from:
 - (1) Chondrocytes
 - (2) Compound epithelial cells
 - (3) Squamous epithelial cells
 - (4) Columnar epithelial cells
- **70.** Montreal protocol was signed in 1987 for control of :
 - (1) Release of Green House gases
 - (2) Disposal of e-wastes
 - (3) Transport of Genetically modified organisms from one country to another
 - (4) Emission of ozone depleting substances
- **71.** Which of the following statements are **true** for the phylum-Chordata?
 - (a) In Urochordata notochord extends from head to tail and it is present throughout their life.
 - (b) In Vertebrata notochord is present during the embryonic period only.
 - (c) Central nervous system is dorsal and hollow.
 - (d) Chordata is divided into 3 subphyla: Hemichordata, Tunicata and Cephalochordata.
 - (1) (a) and (b)
 - (2) (b) and (c)
 - (3) (d) and (c)
 - (4) (c) and (a)
- **72.** Identify the substances having glycosidic bond and peptide bond, respectively in their structure :
 - (1) Cellulose, lecithin
 - (2) Inulin, insulin
 - (3) Chitin, cholesterol
 - (4) Glycerol, trypsin

73. Match the following columns and select the correct option.

Column - I Column - II Placenta (i) Androgens (a) **Human Chorionic** (b) Zona pellucida (ii) Gonadotropin (hCG) (c) Bulbo-urethral (iii) Layer of the ovum glands Leydig cells (d) (iv) Lubrication of the Penis (a) (b) (c) (d) (1) (iii) (ii)(iv) (i) (2)(ii) (iii) (iv) (i) (3)(iv) (iii) (i) (ii) (iii) (4) (i) (iv) (ii)

- 74. If the distance between two consecutive base pairs is 0.34 nm and the total number of base pairs of a DNA double helix in a typical mammalian cell is 6.6×10^9 bp, then the length of the DNA is approximately:
 - (1) 2.2 meters
 - (2) 2.7 meters
 - (3) 2.0 meters
 - (4) 2.5 meters
- **75.** The ovary is half inferior in:
 - (1) Sunflower
 - (2) Plum
 - (3) Brinjal
 - (4) Mustard
- **76.** Identify the **correct** statement with regard to G_1 phase (Gap 1) of interphase.
 - $\begin{array}{c} \hbox{(1)} & \hbox{Cell is metabolically active, grows but does} \\ & \hbox{not replicate its DNA.} \end{array}$
 - (2) Nuclear Division takes place.
 - (3) DNA synthesis or replication takes place.
 - (4) Reorganisation of all cell components takes place.
- 77. Which of the following hormone levels will cause release of ovum (ovulation) from the graffian follicle?
 - (1) Low concentration of LH
 - (2) Low concentration of FSH
 - (3) High concentration of Estrogen
 - (4) High concentration of Progesterone

- **78.** Identify the **correct** statement with reference to human digestive system.
 - (1) Ileum is a highly coiled part.
 - (2) Vermiform appendix arises from duodenum.
 - (3) Ileum opens into small intestine.
 - (4) Serosa is the innermost layer of the alimentary canal.
- **79.** Match the following columns and select the **correct** option.

	Colu	ımn -	I	Column - II	
(a)	Eosii	Eosinophils			Immune response
(b)	Baso	Basophils			Phagocytosis
(c)	Neut	Neutrophils			Release histaminase, destructive
(d)	Lym	Lymphocytes			enzymes Release granules containing histamine
	(a)	(b)	(c)	(d)	
(1)	(i)	(ii)	(iv)	(iii)	
(2)	(ii)	(i)	(iii)	(iv)	
(3)	(iii)	(iv)	(ii)	(i)	
(4)	(iv)	(i)	(ii)	(iii)	

- **80.** The plant parts which consist of two generations one within the other:
 - (a) Pollen grains inside the anther
 - (b) Germinated pollen grain with two male gametes
 - (c) Seed inside the fruit
 - (d) Embryo sac inside the ovule
 - (1) (c) and (d)
 - (2) (a) and (d)
 - (3) (a) only
 - (4) (a), (b) and (c)
- 81. Bt cotton variety that was developed by the introduction of toxin gene of *Bacillus thuringiensis* (Bt) is resistant to:
 - (1) Plant nematodes
 - (2) Insect predators
 - (3) Insect pests
 - (4) Fungal diseases

- **82.** The first phase of translation is:
 - (1) Aminoacylation of tRNA
 - (2) Recognition of an anti-codon
 - (3) Binding of mRNA to ribosome
 - (4) Recognition of DNA molecule
- **83.** Embryological support for evolution was disapproved by:
 - (1) Charles Darwin
 - (2) Oparin
 - (3) Karl Ernst von Baer
 - (4) Alfred Wallace
- 84. Match the following columns and select the correct option.

	Colu	ımn -	I	Column - II	
(a)	6 - 15 pairs of (i) gill slits			(i)	Trygon
(b)	11000	Heterocercal (ii) caudal fin			Cyclostomes
(c)	Air E	Air Bladder			Chondrichthyes
(d)	Poise	Poison sting			Osteichthyes
	(a)	(b)	(c)	(d)	
(1)	(iv)	(ii)	(iii)	(i)	
(2)	(i)	(iv)	(iii)	(ii)	
(3)	(ii)	(iii)	(iv)	(i)	
(4)	(iii)	(iv)	(i)	(ii)	

85. Match the following columns and select the **correct** option.

	Colu	ımn -	I		Column - II
(a)		Clostridium butylicum			Cyclosporin-A
(b)		Trichoderma polysporum			Butyric Acid
(c)		Monascus purpureus			Citric Acid
(d)	Aspe	Aspergillus niger			Blood cholesterol lowering agent
	(a)	(b)	(c)	(d)	
(1)	(i)	(ii)	(iv)	(iii)	
(2)	(iv)	(iii)	(ii)	(i)	
(3)	(iii)	(iv)	(ii)	(i)	
(4)	(ii)	(i)	(iv)	(iii)	

- **86.** Which of the following is **not** an inhibitory substance governing seed dormancy?
 - (1) Phenolic acid
 - (2) Para-ascorbic acid
 - (3) Gibberellic acid
 - (4) Abscisic acid
- 87. Match the following columns and select the **correct** option.

Column - I

Column - II

- (a) Organ of Corti
- $\begin{array}{c} \hbox{(i)} & \quad \text{Connects middle} \\ & \quad \text{ear and pharynx} \end{array}$
- (b) Cochlea
- (ii) Coiled part of the labyrinth
- (c) Eustachian tube
- (iii) Attached to the oval window
- (d) Stapes
- (iv) Located on the basilar membrane
- (a) (b) (c) (d)
- (1) (iv) (ii) (i) (iii)
- (2) (i) (ii) (iv) (iii)
- (3) (ii) (iii) (i) (iv)
- (4) (iii) (i) (iv) (ii)
- **88.** The enzyme enterokinase helps in conversion of:
 - (1) caseinogen into casein
 - (2) pepsinogen into pepsin
 - (3) protein into polypeptides
 - (4) trypsinogen into trypsin
- **89.** Presence of which of the following conditions in urine are indicative of Diabetes Mellitus?
 - (1) Ketonuria and Glycosuria
 - (2) Renal calculi and Hyperglycaemia
 - (3) Uremia and Ketonuria
 - (4) Uremia and Renal Calculi
- **90.** The process responsible for facilitating loss of water in liquid form from the tip of grass blades at night and in early morning is:
 - (1) Imbibition
 - (2) Plasmolysis
 - (3) Transpiration
 - (4) Root pressure

91. A short electric dipole has a dipole moment of 16×10^{-9} C m. The electric potential due to the dipole at a point at a distance of 0.6 m from the centre of the dipole, situated on a line making an angle of 60° with the dipole axis is:

$$\left(\frac{1}{4\pi\epsilon_0} = 9 \times 10^9 \text{ N m}^2/\text{C}^2\right)$$

- (1) 400 V
- (2) zero
- (3) 50 V
- (4) 200 V
- 92. A series LCR circuit is connected to an ac voltage source. When L is removed from the circuit, the phase difference between current and voltage is $\frac{\pi}{3}$. If instead C is removed from the circuit, the phase difference is again $\frac{\pi}{3}$ between current and voltage. The power factor of the circuit is:
 - (1) 1.0
 - (2) -1.0
 - (3) zero
 - (4) 0.5
- 93. Light of frequency 1.5 times the threshold frequency is incident on a photosensitive material. What will be the photoelectric current if the frequency is halved and intensity is doubled?
 - (1) one-fourth
 - (2) zero
 - (3) doubled
 - (4) four times
- **94.** Dimensions of stress are:
 - (1) $[ML^0T^{-2}]$
 - (2) $[ML^{-1}T^{-2}]$
 - (3) $[MLT^{-2}]$
 - (4) $[ML^2T^{-2}]$
- 95. An electron is accelerated from rest through a potential difference of V volt. If the de Broglie wavelength of the electron is 1.227×10^{-2} nm, the potential difference is:
 - (1) $10^3 \, \text{V}$
 - (2) $10^4 \, \text{V}$
 - (3) 10 V
 - (4) $10^2 \,\mathrm{V}$

96. The capacitance of a parallel plate capacitor with air as medium is 6 μF . With the introduction of a dielectric medium, the capacitance becomes 30 μF . The permittivity of the medium is:

$$(\epsilon_0 = 8.85 \times 10^{-12} \text{ C}^2 \text{ N}^{-1} \text{ m}^{-2})$$

- (1) $0.44 \times 10^{-10} \text{ C}^2 \text{ N}^{-1} \text{ m}^{-2}$
- (2) $5.00 \text{ C}^2 \text{ N}^{-1} \text{ m}^{-2}$
- (3) $0.44 \times 10^{-13} \text{ C}^2 \text{ N}^{-1} \text{ m}^{-2}$
- (4) $1.77 \times 10^{-12} \text{ C}^2 \text{ N}^{-1} \text{ m}^{-2}$
- **97.** The solids which have the negative temperature coefficient of resistance are :
 - (1) semiconductors only
 - (2) insulators and semiconductors
 - (3) metals
 - (4) insulators only
- **98.** For transistor action, which of the following statements is **correct**?
 - (1) Both emitter junction as well as the collector junction are forward biased.
 - (2) The base region must be very thin and lightly doped.
 - (3) Base, emitter and collector regions should have same doping concentrations.
 - (4) Base, emitter and collector regions should have same size.
- **99.** A screw gauge has least count of 0.01 mm and there are 50 divisions in its circular scale.


The pitch of the screw gauge is:

- (1) 0.5 mm
- (2) 1.0 mm
- (3) 0.01 mm
- (4) 0.25 mm
- **100.** The phase difference between displacement and acceleration of a particle in a simple harmonic motion is:
 - (1) $\frac{\pi}{2}$ rad
 - (2) zero
 - (3) $\pi \operatorname{rad}$
 - (4) $\frac{3\pi}{2}$ rad

101. A long solenoid of 50 cm length having 100 turns carries a current of 2.5 A. The magnetic field at the centre of the solenoid is:

$$(\mu_0 = 4\pi \times 10^{-7} \text{ T m A}^{-1})$$


- (1) $6.28 \times 10^{-5} \,\mathrm{T}$
- (2) $3.14 \times 10^{-5} \,\mathrm{T}$
- (3) $6.28 \times 10^{-4} \,\mathrm{T}$
- (4) $3.14 \times 10^{-4} \,\mathrm{T}$
- 102. A ball is thrown vertically downward with a velocity of 20 m/s from the top of a tower. It hits the ground after some time with a velocity of 80 m/s. The height of the tower is: $(g=10 \text{ m/s}^2)$
 - (1) 320 m
 - (2) 300 m
 - (3) 360 m
 - (4) 340 m
- **103.** The color code of a resistance is given below:



The values of resistance and tolerance, respectively, are:

- (1) $4.7 \text{ k}\Omega, 5\%$
- (2) $470 \Omega, 5\%$
- (3) $470 \text{ k}\Omega, 5\%$
- (4) $47 \text{ k}\Omega, 10\%$
- **104.** The Brewsters angle i_b for an interface should be :
 - (1) $45^{\circ} < i_b < 90^{\circ}$
 - (2) $i_b = 90^{\circ}$
 - (3) $0^{\circ} < i_b < 30^{\circ}$
 - (4) $30^{\circ} < i_b < 45^{\circ}$
- 105. A ray is incident at an angle of incidence i on one surface of a small angle prism (with angle of prism A) and emerges normally from the opposite surface. If the refractive index of the material of the prism is μ , then the angle of incidence is nearly equal to:
 - (1) μA
 - $(2) \qquad \frac{\mu A}{2}$
 - (3) $\frac{A}{2\mu}$
 - (4) $\frac{2A}{\mu}$

- 106. Two cylinders A and B of equal capacity are connected to each other via a stop cock. A contains an ideal gas at standard temperature and pressure. B is completely evacuated. The entire system is thermally insulated. The stop cock is suddenly opened. The process is:
 - (1) isochoric
 - (2) isobaric
 - (3) isothermal
 - (4) adiabatic
- **107.** For which one of the following, Bohr model is **not** valid?
 - (1) Deuteron atom
 - (2) Singly ionised neon atom (Ne⁺)
 - (3) Hydrogen atom
 - (4) Singly ionised helium atom (He⁺)
- 108. Two bodies of mass 4 kg and 6 kg are tied to the ends of a massless string. The string passes over a pulley which is frictionless (see figure). The acceleration of the system in terms of acceleration due to gravity (g) is:




- (1) g/5
- (2) g/10
- (3) g
- (4) g/2
- **109.** In a certain region of space with volume 0.2 m³, the electric potential is found to be 5 V throughout. The magnitude of electric field in this region is:
 - (1) 1 N/C
 - (2) 5 N/C
 - (3) zero
 - (4) 0.5 N/C

- 110. When a uranium isotope $^{235}_{92}\rm U$ is bombarded with a neutron, it generates $^{89}_{36}\rm Kr$, three neutrons and:
 - (1) $^{101}_{36}$ Kr
 - (2) $^{103}_{36}$ Kr
 - (3) $^{144}_{56}$ Ba
 - (4) $^{91}_{40}$ Zr
- 111. The energy equivalent of 0.5 g of a substance is:
 - (1) $1.5 \times 10^{13} \,\mathrm{J}$
 - (2) $0.5 \times 10^{13} \,\mathrm{J}$
 - (3) $4.5 \times 10^{16} \,\mathrm{J}$
 - (4) $4.5 \times 10^{13} \,\mathrm{J}$
- 112. The mean free path for a gas, with molecular diameter d and number density n can be expressed as:
 - $(1) \qquad \frac{1}{\sqrt{2} \, \operatorname{n}^2 \pi \mathrm{d}^2}$
 - (2) $\sqrt{2} n^2 \pi^2 d^2$
 - (3) $\sqrt{2} n\pi d$
 - $(4) \qquad \frac{1}{\sqrt{2} \, \operatorname{n} \pi \mathrm{d}^2}$
- $\begin{array}{ll} \textbf{113.} & A \ wire \ of \ length \ L, \ area \ of \ cross \ section \ A \ is \ hanging \\ from \ a \ fixed \ support. & The \ length \ of \ the \ wire \\ changes \ to \ L_1 \ when \ mass \ M \ is \ suspended \ from \ its \\ free \ end. & The \ expression \ for \ Young's \ modulus \ is: \end{array}$
 - $(1) \qquad \frac{\text{MgL}}{\text{AL}_1}$
 - $(2) \qquad \frac{MgL}{A(L_1-L)}$
 - $(3) \qquad \frac{\mathrm{MgL}_{1}}{\mathrm{AL}}$
 - $(4) \qquad \frac{\mathrm{Mg(L_1 L)}}{\mathrm{AL}}$
- 114. A spherical conductor of radius 10 cm has a charge of 3.2×10^{-7} C distributed uniformly. What is the magnitude of electric field at a point 15 cm from the centre of the sphere?

$$\left(\frac{1}{4\pi\epsilon_0} = 9 \times 10^9 \ \mathrm{N \ m^2/C^2}\right)$$

- (1) $1.28 \times 10^6 \text{ N/C}$
- (2) $1.28 \times 10^7 \text{ N/C}$
- (3) $1.28 \times 10^4 \text{ N/C}$
- (4) $1.28 \times 10^5 \text{ N/C}$

- 115. The energy required to break one bond in DNA is 10^{-20} J. This value in eV is nearly :
 - (1) 0.06
 - (2) 0.006
 - (3) 6
 - (4) 0.6
- **116.** A body weighs 72 N on the surface of the earth. What is the gravitational force on it, at a height equal to half the radius of the earth?
 - (1) 30 N
 - (2) 24 N
 - (3) 48 N
 - (4) 32 N
- 117. For the logic circuit shown, the truth table is:



- (1) A B Y 0 0 1
 - $0 \quad 1 \quad 1$
 - 1 0 1
- 1 1 0
- (2) A B Y
 - 0 0 1
 - $\begin{array}{cccc} 0 & 1 & 0 \\ 1 & 0 & 0 \end{array}$
- (3) A B Y
 - 0 0 0
 - $\begin{array}{cccc} 0 & 1 & 0 \\ 1 & 0 & 0 \end{array}$
 - 1 1 1
- (4) A B Y
 - 0 0 0
 - 0 1 1
 - $\begin{array}{cccc} 1 & 0 & 1 \\ 1 & 1 & 1 \end{array}$
- 118. In Young's double slit experiment, if the separation between coherent sources is halved and the distance of the screen from the coherent sources is doubled, then the fringe width becomes:
 - (1) four times
 - (2) one-fourth
 - (3) double
 - (4) half

- 119. A capillary tube of radius r is immersed in water and water rises in it to a height h. The mass of the water in the capillary is 5 g. Another capillary tube of radius 2r is immersed in water. The mass of water that will rise in this tube is:
 - (1) 10.0 g
 - (2) 20.0 g
 - (3) 2.5 g
 - (4) 5.0 g
- **120.** A cylinder contains hydrogen gas at pressure of 249 kPa and temperature 27°C.


Its density is: $(R = 8.3 \text{ J mol}^{-1} \text{ K}^{-1})$


- (1) 0.1 kg/m^3
- (2) 0.02 kg/m^3
- (3) 0.5 kg/m^3
- (4) 0.2 kg/m^3
- 121. An iron rod of susceptibility 599 is subjected to a magnetising field of 1200 A m⁻¹. The permeability of the material of the rod is:


$$(\mu_0 = 4\pi \times 10^{-7} \text{ T m A}^{-1})$$


- (1) $2.4\pi \times 10^{-5} \text{ T m A}^{-1}$
- (2) $2.4\pi \times 10^{-7} \text{ T m A}^{-1}$
- (3) $2.4\pi \times 10^{-4} \text{ T m A}^{-1}$
- (4) $8.0 \times 10^{-5} \,\mathrm{T} \,\mathrm{m} \,\mathrm{A}^{-1}$
- 122. Find the torque about the origin when a force of $3\hat{j}$ N acts on a particle whose position vector is $2\hat{k}$ m.
 - (1) $-6\hat{i}$ N m
 - (2) $6\hat{k}$ N m
 - (3) 6i N m
 - (4) $6\hat{j}$ N m
- $\begin{tabular}{ll} \textbf{123.} & The average thermal energy for a mono-atomic gas \\ is: (k_B is Boltzmann constant and T, absolute \\ temperature) \\ \end{tabular}$
 - $(1) \qquad \frac{5}{2} \, \, \mathbf{k_B T}$
 - $(2) \qquad \frac{7}{2} \, \, k_B T$
 - $(3) \qquad \frac{1}{2} \,\, k_B T$
 - (4) $\frac{3}{2} k_B$

- **124.** Assume that light of wavelength 600 nm is coming from a star. The limit of resolution of telescope whose objective has a diameter of 2 m is:
 - (1) $7.32 \times 10^{-7} \, \text{rad}$
 - (2) $6.00 \times 10^{-7} \, \text{rad}$
 - (3) $3.66 \times 10^{-7} \, \text{rad}$
 - (4) $1.83 \times 10^{-7} \, \text{rad}$
- 125. Light with an average flux of 20 W/cm^2 falls on a non-reflecting surface at normal incidence having surface area 20 cm^2 . The energy received by the surface during time span of 1 minute is:
 - (1) $24 \times 10^3 \,\text{J}$
 - (2) $48 \times 10^3 \,\text{J}$
 - (3) $10 \times 10^3 \,\mathrm{J}$
 - (4) $12 \times 10^3 \,\mathrm{J}$
- 126. The ratio of contributions made by the electric field and magnetic field components to the intensity of an electromagnetic wave is: (c = speed of electromagnetic waves)
 - (1) 1:c
 - (2) $1:c^2$
 - (3) c:1
 - (4) 1:1
- 127. Which of the following graph represents the variation of resistivity (ρ) with temperature (T) for copper?









- 128. The quantities of heat required to raise the temperature of two solid copper spheres of radii ${\bf r}_1$ and ${\bf r}_2$ (${\bf r}_1$ = 1.5 ${\bf r}_2$) through 1 K are in the ratio:
 - $(1) \qquad \frac{3}{2}$
 - (2) $\frac{5}{3}$
 - (3) $\frac{27}{8}$
 - (4) $\frac{9}{4}$
- 129. A resistance wire connected in the left gap of a metre bridge balances a 10 Ω resistance in the right gap at a point which divides the bridge wire in the ratio 3:2. If the length of the resistance wire is 1.5 m, then the length of 1 Ω of the resistance wire is:
 - (1) $1.5 \times 10^{-1} \,\mathrm{m}$
 - (2) $1.5 \times 10^{-2} \,\mathrm{m}$
 - (3) $1.0 \times 10^{-2} \,\mathrm{m}$
 - (4) $1.0 \times 10^{-1} \,\mathrm{m}$
- **130.** The increase in the width of the depletion region in a p-n junction diode is due to:
 - (1) both forward bias and reverse bias
 - (2) increase in forward current
 - (3) forward bias only
 - (4) reverse bias only
- 131. A 40 μ F capacitor is connected to a 200 V, 50 Hz ac supply. The rms value of the current in the circuit is, nearly :
 - (1) 2.5 A
 - (2) 25.1 A
 - (3) 1.7 A
 - (4) 2.05 A
- 132. Taking into account of the significant figures, what is the value of 9.99 m 0.0099 m?
 - (1) 9.980 m
 - (2) 9.9 m
 - (3) 9.9801 m
 - (4) 9.98 m

- 133. A charged particle having drift velocity of 7.5×10^{-4} m s⁻¹ in an electric field of 3×10^{-10} Vm⁻¹, has a mobility in m² V⁻¹ s⁻¹ of:
 - (1) 2.5×10^{-6}
 - (2) 2.25×10^{-15}
 - (3) 2.25×10^{15}
 - (4) 2.5×10^6
- 134. In a guitar, two strings A and B made of same material are slightly out of tune and produce beats of frequency 6 Hz. When tension in B is slightly decreased, the beat frequency increases to 7 Hz. If the frequency of A is 530 Hz, the original frequency of B will be:
 - (1) 536 Hz
 - (2) 537 Hz
 - (3) 523 Hz
 - (4) 524 Hz
- 135. Two particles of mass 5 kg and 10 kg respectively are attached to the two ends of a rigid rod of length 1 m with negligible mass.

The centre of mass of the system from the 5 kg particle is nearly at a distance of:

- (1) 67 cm
- (2) 80 cm
- (3) 33 cm
- (4) 50 cm
- **136.** Reaction between benzaldehyde and acetophenone in presence of dilute NaOH is known as:
 - (1) Cross Cannizzaro's reaction
 - (2) Cross Aldol condensation
 - (3) Aldol condensation
 - (4) Cannizzaro's reaction
- **137.** Measuring Zeta potential is useful in determining which property of colloidal solution?
 - (1) Stability of the colloidal particles
 - (2) Size of the colloidal particles
 - (3) Viscosity
 - (4) Solubility

- **138.** A tertiary butyl carbocation is more stable than a secondary butyl carbocation because of which of the following?
 - (1) -R effect of $-CH_3$ groups
 - (2) Hyperconjugation
 - (3) -I effect of $-CH_3$ groups
 - (4) + R effect of CH_3 groups
- **139.** The correct option for free expansion of an ideal gas under adiabatic condition is:
 - (1) $q < 0, \Delta T = 0 \text{ and } w = 0$
 - (2) $q > 0, \Delta T > 0 \text{ and } w > 0$
 - (3) $q = 0, \Delta T = 0 \text{ and } w = 0$
 - (4) $q = 0, \Delta T < 0 \text{ and } w > 0$
- **140.** Match the following:

	Oxide		Nature
(a)	CO	(i)	Basic
(b)	BaO	(ii)	Neutral
(c)	$\mathrm{Al_2O_3}$	(iii)	Acidic
(d)	${\rm Cl_2O_7}$	(iv)	Amphoteric

Which of the following is **correct** option?

	(a)	(b)	(c)	(d)
(1)	(iii)	(iv)	(i)	(ii)
(2)	(iv)	(iii)	(ii)	(i)
(3)	(i)	(ii)	(iii)	(iv)
(4)	(ii)	(i)	(iv)	(iii)

- **141.** Reaction between acetone and methylmagnesium chloride followed by hydrolysis will give:
 - (1) Tert. butyl alcohol
 - (2) Isobutyl alcohol
 - (3) Isopropyl alcohol
 - (4) Sec. butyl alcohol
- 142. The following metal ion activates many enzymes, participates in the oxidation of glucose to produce ATP and with Na, is responsible for the transmission of nerve signals.
 - (1) Calcium
 - (2) Potassium
 - (3) Iron
 - (4) Copper

- **143.** Which of the following is a basic amino acid?
 - (1) Tyrosine
 - (2) Lysine
 - (3) Serine
 - (4) Alanine
- **144.** Identify compound X in the following sequence of reactions:

$$\begin{array}{c} \text{CH}_3 \\ \hline \\ \hline \\ \text{Cl}_2/\text{h}\nu \\ \hline \\ \text{373 K} \\ \end{array}$$

$$(1) \qquad \begin{array}{c} \operatorname{CHCl}_2 \\ \\ \end{array}$$

$$(4) \qquad \begin{array}{c} \operatorname{CH_2Cl} \\ \end{array}$$

145. Which of the following is the **correct** order of increasing field strength of ligands to form coordination compounds?

(1)
$$F^- < SCN^- < C_2O_4^{2-} < CN^-$$

(2)
$$CN^- < C_2O_4^{2-} < SCN^- < F^-$$

(3)
$$SCN^- < F^- < C_2O_4^{2-} < CN^-$$

(4)
$$SCN^- < F^- < CN^- < C_2O_4^{2-}$$


- **146.** Which of the following is a cationic detergent?
 - (1) Cetyltrimethyl ammonium bromide
 - (2) Sodium dodecylbenzene sulphonate
 - (3) Sodium lauryl sulphate
 - (4) Sodium stearate
- **147.** Which one of the followings has maximum number of atoms?
 - (1) $1 \text{ g of } O_2(g) \text{ [Atomic mass of } O = 16]$
 - (2) 1 g of Li(s) [Atomic mass of Li = 7]
 - (3) 1 g of Ag(s) [Atomic mass of Ag = 108]
 - (4) 1 g of Mg(s) [Atomic mass of Mg = 24]
- 148. Identify the incorrect match.

(4)

(b), (ii)

Name **IUPAC Official Name** Unnilunium Mendelevium (a) (i) (b) Unniltrium Lawrencium (ii) (c) Unnilhexium (iii) Seaborgium (d) Unununnium (iv) Darmstadtium (1) (c), (iii) (2)(d), (iv) (3)(a), (i)

149. Which of the following amine will give the carbylamine test?



$$(2) \qquad \begin{array}{c} \text{NHC}_2 \text{H}_5 \\ \\ \end{array}$$

150. Paper chromatography is an example of:

- (1) Thin layer chromatography
- (2) Column chromatography
- (3) Adsorption chromatography
- (4) Partition chromatography

151. A mixture of N_2 and Ar gases in a cylinder contains 7 g of N_2 and 8 g of Ar. If the total pressure of the mixture of the gases in the cylinder is 27 bar, the partial pressure of N_2 is:

[Use atomic masses (in g mol⁻¹): N = 14, Ar = 40]

- (1) 15 bar
- (2) 18 bar
- (3) 9 bar
- (4) 12 bar

- 152. The number of protons, neutrons and electrons in $^{175}_{71}$ Lu, respectively, are:
 - (1) 71, 71 and 104
 - (2) 175, 104 and 71
 - (3) 71, 104 and 71
 - (4) 104, 71 and 71
- **153.** The rate constant for a first order reaction is $4.606 \times 10^{-3} \text{ s}^{-1}$. The time required to reduce 2.0 g of the reactant to 0.2 g is:
 - (1) 500 s
 - (2) 1000 s
 - (3) 100 s
 - (4) 200 s
- **154.** Identify a molecule which does **not** exist.
 - (1) C_2
 - (2) O_2
 - (3) He₂
 - (4) Li₂

155. Hydrolysis of sucrose is given by the following reaction.

Sucrose + $H_2O \rightleftharpoons$ Glucose + Fructose

If the equilibrium constant (K_c) is 2×10^{13} at 300 K, the value of $\Delta_r G^\ominus$ at the same temperature will be :

- (1) $8.314 \,\mathrm{J}\,\mathrm{mol}^{-1}\mathrm{K}^{-1} \times 300 \,\mathrm{K} \times \ln(3 \times 10^{13})$
- (2) $-8.314 \,\mathrm{J}\,\mathrm{mol}^{-1}\mathrm{K}^{-1} \times 300 \,\mathrm{K} \times \ln(4 \times 10^{13})$
- (3) $-8.314 \,\mathrm{J}\,\mathrm{mol}^{-1}\mathrm{K}^{-1} \times 300 \,\mathrm{K} \times \ln(2 \times 10^{13})$
- (4) $8.314 \,\mathrm{J}\,\mathrm{mol}^{-1}\mathrm{K}^{-1} \times 300 \,\mathrm{K} \times \ln(2 \times 10^{13})$
- 156. For the reaction, $2Cl(g) \to Cl_2(g),$ the $\boldsymbol{correct}$ option is :
 - (1) $\Delta_r H < 0 \text{ and } \Delta_r S > 0$
 - (2) $\Delta_r H < 0$ and $\Delta_r S < 0$
 - (3) $\Delta_r H > 0$ and $\Delta_r S > 0$
 - (4) $\Delta_r H > 0$ and $\Delta_r S < 0$
- 157. Find out the solubility of Ni(OH) $_2$ in 0.1 M NaOH. Given that the ionic product of Ni(OH) $_2$ is 2×10^{-15} .
 - (1) $1 \times 10^{-13} \,\mathrm{M}$
 - (2) $1 \times 10^8 \,\mathrm{M}$
 - (3) $2 \times 10^{-13} \,\mathrm{M}$
 - (4) $2 \times 10^{-8} \,\mathrm{M}$

- **158.** On electrolysis of dil.sulphuric acid using Platinum (Pt) electrode, the product obtained at anode will be:
 - (1) H_2S gas
 - (2) SO_2 gas
 - (3) Hydrogen gas
 - (4) Oxygen gas
- **159.** Which of the following is **not** correct about carbon monoxide?
 - (1) The carboxyhaemoglobin (haemoglobin bound to CO) is less stable than oxyhaemoglobin.
 - (2) It is produced due to incomplete combustion.
 - (3) It forms carboxyhaemoglobin.
 - (4) It reduces oxygen carrying ability of blood.
- 160. The number of Faradays(F) required to produce 20 g of calcium from molten $CaCl_2$ (Atomic mass of Ca = 40 g mol⁻¹) is:
 - (1) 3
 - (2) 4
 - (3) 1
 - (4) 2
- **161.** Elimination reaction of 2-Bromo-pentane to form pent-2-ene is:
 - (a) β-Elimination reaction
 - (b) Follows Zaitsev rule
 - (c) Dehydrohalogenation reaction
 - (d) Dehydration reaction
 - (1) (b), (c), (d)
 - (2) (a), (b), (d)
 - (3) (a), (b), (c)
 - (4) (a), (c), (d)
- **162.** What is the change in oxidation number of carbon in the following reaction?

$$CH_4(g) + 4Cl_2(g) \rightarrow CCl_4(l) + 4HCl(g)$$

- (1) -4 to +4
- (2) 0 to -4
- (3) + 4 to + 4
- (4) 0 to + 4

- **163.** Which of the following alkane cannot be made in good yield by Wurtz reaction?
 - (1) n-Heptane
 - (2) n-Butane
 - (3) n-Hexane
 - (4) 2,3-Dimethylbutane
- 164. Sucrose on hydrolysis gives:
 - (1) α -D-Glucose + β -D-Fructose
 - (2) α -D-Fructose + β -D-Fructose
 - (3) β -D-Glucose + α -D-Fructose
 - (4) α -D-Glucose + β -D-Glucose
- **165.** Identify the **incorrect** statement.
 - (1) Interstitial compounds are those that are formed when small atoms like H, C or N are trapped inside the crystal lattices of metals.
 - (2) The oxidation states of chromium in ${\rm CrO}_4^{2-}$ and ${\rm Cr}_2{\rm O}_7^{2-}$ are not the same.
 - (3) ${\rm Cr}^{2+}(d^4)$ is a stronger reducing agent than ${\rm Fe}^{2+}(d^6)$ in water.
 - (4) The transition metals and their compounds are known for their catalytic activity due to their ability to adopt multiple oxidation states and to form complexes.
- **166.** HCl was passed through a solution of CaCl₂, MgCl₂ and NaCl. Which of the following compound(s) crystallise(s)?
 - (1) Only MgCl₂
 - (2) NaCl, MgCl₂ and CaCl₂
 - (3) Both MgCl₂ and CaCl₂
 - (4) Only NaCl

- 167. Identify the correct statements from the following:
 - $CO_{9}(g)$ is used as refrigerant for ice-cream (a) and frozen food.
 - The structure of C_{60} contains twelve six (b) carbon rings and twenty five carbon rings.
 - ZSM-5, a type of zeolite, is used to convert (c) alcohols into gasoline.
 - (d) CO is colorless and odourless gas.
 - (1) (b) and (c) only
 - (2)(c) and (d) only
 - (3)(a), (b) and (c) only
 - (4)(a) and (c) only
- 168. An increase in the concentration of the reactants of a reaction leads to change in:
 - (1) threshold energy
 - (2)collision frequency
 - (3)activation energy
 - (4) heat of reaction
- The calculated spin only magnetic moment of Cr^{2+} 169. ion is:
 - (1) $5.92\,\mathrm{BM}$
 - (2) $2.84\,\mathrm{BM}$
 - (3) $3.87\,\mathrm{BM}$
 - (4) $4.90\,\mathrm{BM}$
- 170. Match the following and identify the correct option.
 - $CO(g) + H_2(g)$ (a)
- $Mg(HCO_3)_2 +$ (i) Ca(HCO₃)₂
- (b) Temporary hardness of water
- (ii) An electron deficient hydride
- (c) B_2H_6
- (iii) Synthesis gas
- (d) H_2O_2
- Non-planar (iv) structure
- (a) (b) **(c)** (d)
- (1)(iii)
 - (iv)

(i)

(ii)


- (ii) (i)
- (2)(i)
- (iii)
- (ii) (iv)
- (3)(iii)
- (ii) (iv)
- (4)(iii)
- (i) (iv)
- The mixture which shows positive deviation from Raoult's law is:
 - (1) Acetone + Chloroform
 - (2)Chloroethane + Bromoethane
 - (3)Ethanol + Acetone
 - Benzene + Toluene (4)

172. Anisole on cleavage with HI gives:

$$(1) \hspace{1cm} \begin{array}{c} \text{OH} \\ \\ \\ \end{array} + \text{C}_2 \text{H}_5 \text{I} \\ \end{array}$$

$$(2) \hspace{1cm} + C_2 H_5 O H$$

(3)
$$OH$$
 $+ CH_3I$



- Urea reacts with water to form A which will decompose to form B. B when passed through Cu²⁺ (aq), deep blue colour solution C is formed. What is the formula of **C** from the following?
 - $Cu(OH)_2$ (1)
 - (2) $CuCO_3 \cdot Cu(OH)_2$
 - (3) $CuSO_4$
 - $[Cu(NH_2)_4]^{2+}$ (4)
- The freezing point depression constant $(\boldsymbol{K}_{\!f}\!)$ of benzene is $5.12 \text{ K kg mol}^{-1}$. The freezing point depression for the solution of molality 0.078 m containing a non-electrolyte solute in benzene is (rounded off up to two decimal places):
 - $0.40\,\mathrm{K}$ (1)
 - (2) $0.60\,\mathrm{K}$
 - (3) $0.20\,\mathrm{K}$
 - (4) $0.80\,\mathrm{K}$

- **175.** Which of the following oxoacid of sulphur has -O-O- linkage?
 - (1) $H_2S_2O_8$, peroxodisulphuric acid
 - (2) $H_2S_2O_7$, pyrosulphuric acid
 - (3) H_2SO_3 , sulphurous acid
 - (4) H_2SO_4 , sulphuric acid
- **176.** Identify the **correct** statement from the following:
 - (1) Vapour phase refining is carried out for Nickel by Van Arkel method.
 - (2) Pig iron can be moulded into a variety of shapes.
 - (3) Wrought iron is impure iron with 4% carbon.
 - ${\rm (4)} \qquad {\rm Blister\ copper\ has\ blistered\ appearance\ due} \\ {\rm to\ evolution\ of\ CO_2}.$
- **177.** Which of the following is a natural polymer?
 - (1) polybutadiene
 - (2) poly (Butadiene-acrylonitrile)
 - (3) *cis*-1,4-polyisoprene
 - (4) poly (Butadiene-styrene)
- 178. An element has a body centered cubic (bcc) structure with a cell edge of 288 pm. The atomic radius is:

$$(1) \qquad \frac{4}{\sqrt{3}} \times 288 \text{ pm}$$

(2)
$$\frac{4}{\sqrt{2}} \times 288 \text{ pm}$$

$$(3) \qquad \frac{\sqrt{3}}{4} \times 288 \text{ pm}$$


(4)
$$\frac{\sqrt{2}}{4} \times 288 \text{ pm}$$

179. An alkene on ozonolysis gives methanal as one of the product. Its structure is:

$$CH_2-CH=CH_2$$

$$\begin{array}{c} \operatorname{CH_2CH_2CH_3} \\ \\ \end{array} \tag{2}$$

$$CH = CH - CH_3$$
(3)



- **180.** Which of the following set of molecules will have zero dipole moment?
 - (1) Nitrogen trifluoride, beryllium difluoride, water, 1,3-dichlorobenzene
 - (2) Boron trifluoride, beryllium difluoride, carbon dioxide, 1,4-dichlorobenzene
 - (3) Ammonia, beryllium difluoride, water, 1,4-dichlorobenzene
 - (4) Boron trifluoride, hydrogen fluoride, carbon dioxide, 1,3-dichlorobenzene

- o 0 o -