

05/08/2022

Slot-2



Corporate Office : Aakash Tower, 8, Pusa Road, New Delhi-110005 | Ph.: 011-47623456

## Answers & Solutions

Time : 45 min.

M.M. : 200

### *for* CUET UG-2022 (Physics)

#### IMPORTANT INSTRUCTIONS:

1. The test is of 45 Minutes duration.
2. The test contains 50 Questions out of which 40 questions need to be attempted.
3. Marking Scheme of the test:
  - a. Correct answer or the most appropriate answer: Five marks (+5)
  - b. Any incorrect option marked will be given minus one mark (–1).
  - c. Unanswered/Marked for Review will be given no mark (0).

Choose the correct answer :

Question ID: 102001

During the formation of depletion layer around the imaginary pn-junction, electrons diffuse from n-region to p-region because.

- (A) Electron concentration in n-segment is higher than electron concentration in p-segment.
- (B) Holes of p-segment attract the electrons
- (C) Electrons are lighter and have higher drift velocity
- (D) Hole concentration in n-segment is higher than the electron concentration in p-segment

Answer (A)

**Sol.** During the formation of depletion layer around the imaginary pn-junction, electron diffuses from n-region to p-region because electron concentration in n-segment is higher than electron concentration in p-segment.

Question ID: 102002

The most exotic diamagnetic materials are super conductors. They exhibit perfect diamagnetism and perfect conductivity. The values of magnetic susceptibility ( $\chi$ ) and relative permeability of such materials have values

- (A)  $\chi = 1$ ,  $\mu_r = 0$
- (B)  $\chi = -1$ ,  $\mu_r = 0$
- (C)  $\chi = 0$ ,  $\mu_r = 1$
- (D)  $\chi = -1$ ,  $\mu_r = 1$

Answer (B)

**Sol.** Relation between susceptibility ( $\chi$ ) and relative permeability ( $\mu_r$ )

$$\mu_r = \chi + 1$$

If diamagnetic material behave like

super conductor  $\mu_r = 0$


$$0 = \chi + 1$$

$$\chi = -1$$

Question ID: 102003

The variation of number of undecayed nuclei with time is best represented by

(Where  $N_0$ -initial number of atoms)




Answer (B)

$$\text{Sol. } N = N_0 e^{-\lambda t}$$


$N$  is the number of undecayed nuclei at any time  $t$

$N_0$  is the number of nuclei at  $t = 0$



Question ID: 102004

Which one of the following correctly represent the variation of electric field and electric potential with a distance  $r$  from a point charge?




Answer (C)

**Sol.** Variation of electric field ( $E$ ) of point charge(s) with


$$\text{the distance } (r) \text{ is } E = \frac{kq}{r^2} \Rightarrow E \propto \frac{1}{r^2}$$

Variation of electric potential ( $V$ ) of point charge ( $q$ )

$$\text{with distance } (r) \text{ } V = \frac{kq}{r} \Rightarrow V \propto \frac{1}{r}$$




Question ID: 102005



The above circuit potential difference between point A and B ( $V_A - V_B$ ) =

- (A)  $\frac{4}{3}V$  (B)  $\frac{2}{3}V$   
(C)  $\frac{1}{3}V$  (D)  $\frac{5}{3}V$

Answer (B)



$$R_{\text{eff}} = 4 + \frac{3 \times 6}{9} = 6 \Omega$$

$$I = \frac{V}{R_{\text{eff}}} = \frac{12}{6} = 2 \text{ A}$$

K.V.L in loop ... (1)

$$12 - 4 \times 2 - I_1 \times 3 = 0$$

$$I_1 = \frac{4}{3} \text{ A}$$

K.V.L in loop ... (2)

$$12 - 4 \times 2 - 6I_2 = 0$$

$$I_2 = \frac{2}{3} \text{ A}$$

$$V_0 - V_A = 1 \times \frac{4}{3}$$

$$V_0 - V_A = \frac{4}{3} \quad \dots (i)$$

$$V_0 - V_B = \frac{2}{3} \times 3$$

$$V_0 - V_B = 2V \quad \dots (ii)$$

$$(ii) - (i)$$

$$V_A - V_B = 2 - \frac{4}{3} = \frac{2}{3}V$$

Question ID: 102006

A circular lamina of radius ' $R$ ' is having surface charge density  $\sigma \text{ cm}^{-2}$ . Electric field at axial distance ' $2R$ ' is

- (A)  $\frac{\sigma}{4\pi\epsilon_0 R^2}$  (B)  $\frac{\sigma}{\epsilon_0}$   
(C)  $\frac{\sigma}{2\epsilon_0}$  (D)  $\frac{\sigma}{4\epsilon_0}$


Answer (NA)

**Sol.**  $E = \frac{\sigma}{2\epsilon_0} (1 - \cos\phi)$ , where  $\cos\phi = \frac{2R}{\sqrt{4R^2 + R^2}}$

$$E = \frac{\sigma}{2\epsilon_0} \left( 1 - \frac{2}{\sqrt{5}} \right)$$

Question ID: 102007

Two point charges  $+5 \mu\text{C}$  and  $-5 \mu\text{C}$  are placed at O (0 mm, 0 mm) and P(3 mm, 4 mm) respectively force on  $+5 \mu\text{C}$  is



(A)  $1.8(3\hat{i} - 4\hat{j}) \times 10^{-3} \text{ N}$

(B)  $-1.8(3\hat{i} + 4\hat{j}) \times 10^{-3} \text{ N}$

(C)  $1.8(3\hat{i} + 4\hat{j}) \times 10^{-3} \text{ N}$

(D)  $1.8(-3\hat{i} + 4\hat{j}) \times 10^{-3} \text{ N}$

Answer (C)

**Sol.** Force between two point charge  $q_1$  and  $q_2$

$$\vec{F} = \frac{kq_1q_2(\vec{r}_2 - \vec{r}_1)}{|\vec{r}_2 - \vec{r}_1|^3}$$

$$\vec{F} = \frac{9 \times 10^9 \times 5 \times 5 \times 10^{-12} (4\hat{j} + 3\hat{i}) \times 10^{-3}}{(\sqrt{4^2 + 3^2}) \times 10^{-9}}$$

$$\vec{F} = 1.8(3\hat{i} + 4\hat{j}) \times 10^{-3} \text{ N}$$

**Question ID: 102008**

Which of the following statements is true for a  $p$ -type semiconductor?


- (A) Holes are minority carries and pentavalent atoms are the dopant
- (B) Electrons are minority carries and pentavalent atoms are dopant
- (C) Holes are majority carriers and trivalent atoms are the dopant
- (D) Electrons are majority carriers and trivalent atoms are the dopant

**Answer (C)**

**Sol.** In  $p$ -type semiconductor holes are majority carrier and trivalent atom are dopant.

**Question ID: 102009**

A graph for variation of photo current with anode potential is given for different intensities of incident radiation:




Choose the correct order of anode intensities.

- (A)  $I_1 > I_2 > I_3$
- (B)  $I_1 = I_2 = I_3$
- (C)  $I_1 < I_2 < I_3$
- (D)  $I_1 < I_2 > I_3$

**Answer (C)**

**Sol.** Photo current is directly proportional to intensity of incident light.



**Question ID: 102010**

NPN transistor is set up as amplifier in common emitter mode, output signal voltage obtained across the load resistance of  $4 \text{ k}\Omega$  is 1 volt. Calculate input signal voltage. Assume current amplification fevelar = 250 and base resistance =  $1 \text{ k}\Omega$ :

- (A) 4 mV
- (B) 2.5 mV
- (C) 1 mV
- (D) Insufficient data

**Answer (C)**

$$\text{Sol. } I_c = \frac{V_c}{R_c} = \frac{1}{4 \times 10^3} = 0.25 \text{ mA}$$

$$I_B = \frac{I_c}{\beta} = \frac{0.25 \times 10^{-3}}{250} = 10^{-6} \text{ A}$$

$$V_i = I_B R_B = 10^{-6} \times 10^3 = 1 \text{ mV}$$

**Question ID: 102011**

Distance of ray optics as a good approximation, when the aperture is 5 mm wide and the wavelength is 600 nm is nearly :

- (A) 18 m
- (B) 15 m
- (C) 4.43 m
- (D) 8.34 m

**Answer (NA)**


$$\text{Sol. Fresnel distance } (Z) = \frac{a^2}{\lambda}$$

Here  $a$  is aperture width  $\lambda$  is wavelength of light

$$Z = \frac{(5 \times 10^{-3})^2}{600 \times 10^{-9}} = \frac{25}{6} \times 10 \approx 41.67 \text{ m}$$

**Question ID: 102012**


Three charges  $Q$ ,  $+q$  and  $+q$  are placed at the vertices of an equilateral triangle of side  $l$ , as shown in the figure. If the net electrostatic energy of the system is zero, the  $Q$  is equal to:



- (A)  $+\frac{q}{2}$
- (B) Zero
- (C)  $-\frac{q}{2}$
- (D)  $-q$

**Answer (C)**

Sol.



$$U = \frac{kq^2}{l} + \frac{kqQ}{l} + \frac{kqQ}{l}$$

Given,  $U = 0$ 

$$= \frac{kq^2}{l} + \frac{2kqQ}{l} = 0$$

$$Q = \frac{-q}{2}$$

Question ID: 102013

If the ratio of maximum to minimum resultant intensity in interference pattern of two waves is 25 : 4, then, ratio of amplitudes two waves is :

- (A) 5 : 2  
(B) 2 : 5  
(C) 7 : 3  
(D) 49 : 9

Answer (C)

$$\text{Sol. } \frac{I_{\max}}{I_{\min}} = \frac{(\sqrt{I_1} + \sqrt{I_2})^2}{(\sqrt{I_1} - \sqrt{I_2})^2}$$

$$\frac{25}{4} = \left( \frac{\sqrt{I_1} + \sqrt{I_2}}{\sqrt{I_1} - \sqrt{I_2}} \right)^2$$

$$\frac{5}{2} = \left( \frac{\sqrt{I_1} + \sqrt{I_2}}{\sqrt{I_1} - \sqrt{I_2}} \right)$$

$$3\sqrt{I_1} = 7\sqrt{I_2} \Rightarrow \frac{I_1}{I_2} = \frac{49}{9}$$


$$I \propto A^2$$

$$\Rightarrow A \propto \sqrt{I}$$

$$\Rightarrow \frac{A_1}{A_2} = \sqrt{\frac{I_1}{I_2}} = \sqrt{\frac{49}{9}} = \frac{7}{3}$$

Question ID: 102014

In the circuit given below the current is to be measured. If the ammeter shown is a galvanometer with a resistance  $R_G = 60 \Omega$  but converted to an ammeter by a shunt resistance  $r_s = 0.02 \Omega$ , then the flow of current is nearly :




- (1) 0.5 A                      (2) 3.02 A  
(3) 0.99 A                    (4) 1.5 A  
(A) 1                            (B) 2  
(C) 3                            (D) 4

Answer (NA)

Sol. Emf of battery is not given Data is insufficient

Question ID: 102015

Consider a capacitor circuit charged by a battery to a potential difference  $V_0$  as shown in the figure.




When a dielectric slab is inserted into both the capacitors, new potential difference will be

- (A)  $V_0$                       (B)  $\frac{V_0}{K}$   
(C)  $\frac{8V_0}{5K}$                     (D)  $\frac{8V_0}{K}$

Answer (C)

Sol.



On inserting dielectric charge on the combination of capacitor remains same  $q = cV$

$$4C_0V_0 = \left(KC_0 + \frac{3C_0K}{2}\right)V_1$$

$$V_1 = \frac{8V_0}{5K}$$

**Question ID:102016**

An intrinsic semiconductor has  $5 \times 10^{28}$  atoms/m<sup>3</sup>. It is doped by 0.01 ppm. concentration of arsenic. If  $n_i = 1.5 \times 10^{16}$  m<sup>3</sup>, then number of holes in the  $n$ -type semiconductor will be

- (A)  $5 \times 10^{28}/\text{m}^3$  (B)  $5 \times 10^{11}/\text{m}^3$   
(C)  $3.0 \times 10^{11}/\text{m}^3$  (D)  $4.5 \times 10^{11}/\text{m}^3$

**Answer (D)**

**Sol.** 0.01 atom of Si doped out of  $10^6$  atom 1 ppm.

$$\begin{aligned} \text{In } 5 \times 10^{28} \text{ atom number of doped} &= \frac{5 \times 10^{28}}{\frac{10^6}{0.01}} \\ &= 5 \times 10^{20} \end{aligned}$$

1 as atom creates  $1e^-$  excess so number of excess electron =  $5 \times 10^{20} = n_e$

$$\text{also } n_e n_h = n_i^2$$

$$\begin{aligned} n_h &= \frac{n_i^2}{n_e} = \frac{(1.5 \times 10^{16})^2}{5 \times 10^{20}} \\ &= 0.45 \times 10^{12} \\ &= 4.5 \times 10^{11}/\text{m}^3 \end{aligned}$$

**Question ID:102017**

In the block diagram of a simple modulator, a square law device generates DC and sinusoidal of frequencies  $\omega_m$ ,  $2\omega_m$ ,  $\omega_c$ ,  $2\omega_c$ ,  $\omega_c - \omega_m$  and  $\omega_c + \omega_m$ . The device which rejects dc and unwanted frequencies and retains frequencies,  $\omega_c$ ,  $\omega_c - \omega_m$  and  $\omega_c + \omega_m$  is called

- (A) Envelope detector (B) Band pass filter  
(C) Rectifier (D) Amplifier

**Answer (B)**

**Sol.** The device which reject dc and unwanted frequencies and retains frequencies,  $\omega_c$ ,  $\omega_c - \omega_m$  and  $\omega_c + \omega_m$  is called Band pass filter

**Question ID:102018**

An expression for the frequency of revolution of the electron in the Bohr's orbit is :

- (A)  $\frac{me^4}{4\epsilon_0^2 h^3} \times \frac{1}{n^3}$  (B)  $\frac{me^4}{8\epsilon_0^2 h^2} \times \frac{1}{n^3}$   
(C)  $\frac{me^4}{4\epsilon_0^2 ch^3} \times \frac{1}{n^3}$  (D)  $\frac{me^4}{8\epsilon_0^2 ch^2} \times \frac{1}{n^3}$

**Answer (A)**

**Sol.** In Bohr's orbit

$$mvr = \frac{nh}{2\pi} \quad \dots(i)$$

$$\frac{e^2 z}{4\pi\epsilon_0 r^2} = \frac{mv^2}{r}$$

$$mv^2 r = \frac{e^2 z}{4\pi\epsilon_0} \quad \dots(ii)$$

On putting the value of  $mvr$  from equation (i)

$$v \times \frac{nh}{2\pi} = \frac{e^2 z}{4\pi\epsilon_0}$$

$$v = \left( \frac{e^2 z}{2nh} \right)$$

$$\frac{m \times e^2 z}{2nh} r = \frac{nh}{2\pi}$$

$$r = \frac{n^2 h^2}{\pi m e^2 z}$$

$$\text{Now, } f = \frac{v}{2\pi r}$$


$$\begin{aligned} &= \frac{e^2 z}{2nh} \times \frac{2\pi n^2 h^2}{\pi m e^2 z} \\ &= \frac{me^4 z^2}{4\epsilon_0^2 h^3} \times \frac{1}{n^3} \end{aligned}$$

**Question ID:102019**

Current in an alternating circuit is ahead of voltage by a phase difference of  $\frac{\pi}{3}$ . The alternating circuit is

- (A) C circuit  
(B) LC circuit  
(C) LR circuit  
(D) L circuit

**Answer (NA)**



Current lead voltage by  $\frac{\pi}{3}$  therefore alternating circuit will be RC. No option is correct.

**Question ID:102020**

A particle moves in a closed orbit around the origin, due to a force which is directed towards the origin. The de-Broglie wavelength of particle varies cyclically between two values ( $\lambda_1$ ), ( $\lambda_2$ ) with ( $\lambda_1 > \lambda_2$ ) then, the correct statement from the following are:

- (a) The particle could be moving in a circular orbit with origin as centre
- (b) The particle could be moving in an elliptical orbit with origin as its focus.
- (c) When the de-Broglie wave length is  $\lambda_1$ , the particle is nearer to origin then when its value is  $\lambda_2$ .
- (d) When the de Broglie wavelength is  $\lambda_2$ , the particle is nearer to the origin than when its value is  $\lambda_1$ .

- (A) (a) and (d) only      (B) (b) and (d) only
- (C) (b) and (c) only      (D) (c) and (a) only

**Answer (B)**


Sol. de-Broglie wavelength of a moving particle  $\lambda = \frac{h}{mv}$

$\lambda_1 > \lambda_2$  it means  $v_1 > v_2$  velocity of the particle will be greater at nearer point on a elliptical orbit.

**Question ID: 102021**

A circular coil made up of copper, is placed on a table top. Magnetic field is normally outwards. Now, the magnetic field is gradually decrease. The direction of induced current as seen from the top side is

- (A) Clock wise
- (B) Anti clock wise
- (C) Current will not be induced
- (D) Information is insufficient

**Answer (B)**


Magnetic field increasing outwards gradually. As per Lenz's law current will flow anticlockwise so that outward flux is supported.

**Question ID: 102022**

In case of Bohr's model of atoms, which of the following statement is incorrect?


- (A) Bohr's model is applicable only for one electron atom/
- (B) Bohr's model correctly predicts the frequencies of light emitted by one electron atoms/ions.
- (C) Bohr's model correctly predicts the intensity variations in the light of different frequencies emitted by one-electron atoms, ions
- (D) Bohr's model does not consider the electron-electron interaction in an atom

**Answer (C)**

Sol. Bohr's model correctly predicts the frequencies of light emitted by one electron atom/ions and not the intensity variation in the light of different frequencies of emitted by one-electron atoms, ion

**Question ID: 102023**

Current through the galvanometer in the circuit given below is



- (A) 6.13 mA      (B) 33.9 mA
- (C) 6.69 mA      (D) 51.7 mA

**Answer (NA)**

Sol. As A and C is shorted, no current will flow through galvanometer




**Question ID: 102024**

During the process of Beta decay the ratio of neutron to proton:

- (A) increases
- (B) decreases
- (C) remain unchanged
- (D) first increases and then decreases

**Answer (B)**


**Sol.** Neutron to proton ratio of an atom nucleus is the ratio of the number of neutrons to its number of proton. This ratio generally increases with increase in atomic number in stable nuclei. Beta decay decreases the ratio of neutron to proton eq.



It is clearly visible that the neutron proton ratio decreases in Beta decay


**Question ID: 102025**

The resultant magnetic field at point O, due to current I in the conductor shown in figure is



- (A)  $\vec{B} = \frac{\mu_0 I}{4r} \left[ \frac{1}{\pi} + 1 \right]$
- (B)  $\vec{B} = \frac{\mu_0 I}{4r} \left[ \frac{2}{\pi} + 1 \right]$
- (C)  $\vec{B} = \frac{\mu_0 I}{4r} \left[ \frac{2}{\pi} - 1 \right]$
- (D)  $\vec{B} = \frac{\mu_0 I}{4r} \left[ \frac{1}{\pi} - 1 \right]$

**Answer (B)**



$$\vec{B}_1 = \frac{\mu_0 I}{4\pi\gamma} (-\hat{k})$$

$$\vec{B}_2 = \frac{\mu_0 I}{4\pi} (\hat{k})$$

$$\vec{B}_3 = \frac{\mu_0 I}{4\pi} (-\hat{k})$$

$$\vec{B} = \vec{B}_1 + \vec{B}_2 + \vec{B}_3$$

$$= \left( \frac{\mu_0 I}{4\pi\gamma} + \frac{\mu_0 I}{4\gamma} + \frac{\mu_0 I}{4\pi\gamma} \right) = \frac{\mu_0 I}{4\gamma} \left( \frac{2}{\pi} + 1 \right) (-\hat{k})$$


**Question ID: 102026**

Correct symbol of NPN transistor is :

- (A)
- (B)
- (C)
- (D)

**Answer (A)**

**Sol.** Symbol of NPN transistor



**Question ID: 102027**

In a typical set-up of photo electric effect, the original source of blue light is replaced with a source of red light of testy. Which of the following phenomenon is observed?

- (A) Photoelectric current increases
- (B) Photoelectric current decreases
- (C) Stopping potential increases
- (D) Stopping potential decreases

**Answer (D)**

**Sol.** Frequency of red light is less than frequency of blue light therefore kinetic energy of emitted electron decreases and stopping potential decreases.



**Question ID: 102028**

A cycle wheel is rotated with a speed of 240 rpm in a plane normal to horizontal component of earth's magnetic field of 0.36 G. Find the value of induced current between the rim and axle of the wheel given, there are 16 spokes, each 70 cm long. The resistance of each spoke is  $2\ \Omega$ .

- (A)  $1.76 \times 10^{-1}\text{ A}$  (B)  $1.76 \times 10^{-3}\text{ A}$   
 (C)  $1.10 \times 10^{-5}\text{ A}$  (D)  $1.10 \times 10^{-4}\text{ A}$

**Answer (B\*)**

**Sol.** Induced emf ( $\mathcal{E}$ ) =  $\frac{\beta\omega l^2}{2}$

$$= \frac{0.36 \times 10^{-4} \times 2\pi f \times (70 \times 10^{-2})^2}{2}$$

$$= \frac{36 \times 10^{-6} \times 8\pi \times 4900 \times 10^{-4}}{2}$$

$$= 18 \times 8 \times 6.28 \times 49 \times 10^{-8}$$

$$R_{\text{eff}} = \frac{2}{16} = \frac{1}{8}$$

$$I = R_{\text{eff}} = \frac{18 \times 8 \times 6.28 \times 49 \times 10^{-8}}{\frac{1}{8}}$$

$$\approx 1.76 \times 10^{-3}\text{ A}$$

**Question ID: 102029**

Electron is accelerated by 100 volt potential difference. Its wavelength will be comparable to that of :

- (A) Infrared radiator (B) Microwave  
 (C) X-rays (D) Radio waves

**Answer (C)**

**Sol.** de-Broglie wavelength of electron =  $\frac{1.227}{\sqrt{V}}\text{ nm}$

$$= \frac{1.227}{\sqrt{100}}$$

$$= 0.1227\text{ nm}$$

Range of X-rays (0.01 – 10) nm

**Question ID: 102030**

Superconductors show :

- (A) Perfect diamagnetism  
 (B) Perfect paramagnetic  
 (C) Perfect ferromagnetism  
 (D) Perfect anti ferromagnetism

**Answer (A)**

**Sol.** Superconductor show perfect diamagnetism.

**Question ID: 102031**

Two coils have a mutual inductance of 0.005 H. The current changes in the first coil according to the equation  $I = I_0 \sin \omega t$ , where  $I_0 = 10\text{ A}$  and  $\omega = 100\pi\text{ rad/s}$ .

The maximum value of emf in the second coil is :

- (A)  $2\pi$  (B)  $5\pi$   
 (C)  $6\pi$  (D)  $12\pi$

**Answer (B)**

**Sol.** emf induced in the second coil ( $\mathcal{E}_s$ )

$$\mathcal{E}_s = -L \frac{di_f}{dt}$$

$$\Rightarrow |\mathcal{E}_s| = L \frac{di_s}{dt}$$

$$= L \frac{d}{dt}(I_0 \sin \omega t)$$

$$= L I_0 \omega \cos \omega t$$

$$\Rightarrow |\mathcal{E}_s|_{\text{max}} = L I_0 \omega$$

$$= 0.005 \times 10 \times 100\pi$$

$$= 5\pi$$

**Question ID: 102032**

Bohr model of atom is valid for :

- (A) Only hydrogen atom  
 (B) Only one electron atom  
 (C) Only one electron atoms and ions  
 (D) All the atoms

**Answer (C)**

**Sol.** Bohr's model is valid only for the single electronic system or only one electron atoms or ions.

**Question ID: 102033**

An ac source of emf  $V(t) = V_0 \sin \omega t$  is put across a pure capacitor. The value of angular frequency of instantaneous power is:

- (A) 0 (B)  $\omega$   
 (C)  $2\omega$  (D)  $\frac{\omega}{2}$

**Answer (C)**

**Sol.**  $V(t) = V_0 \sin \omega t$

In a capacitor  $I(t) = I_0 \sin\left(\omega t + \frac{\pi}{2}\right)$

$$= I_0 \cot \omega t$$

$$P_{\text{inf}} = V(t)I(t)$$

$$= V_0 \sin \omega t I_0 \cos \omega t$$

$$= \frac{V_0 I_0}{3} \sin 2\omega t$$

Angular frequency =  $2\omega$

**Question ID: 102034**

How does voltage sensitivity of a galvanometer will change as number of turns of its coil are doubled?

- (A) Halved
- (B) Doubled
- (C) Remains same
- (D) Insufficient information

**Answer (C)**

**Sol.**  $\frac{\phi}{V} = \frac{NBA}{R}$

As number of turns is doubled, Resistance  $R$  will also get doubled. Hence voltage sensitivity remains same.

**Question ID: 102035**

The Brewster's angle for air to glass interface then an unpolarised light is incident on a glass ( $n = 1.5$ ), such that reflected and refracted rays are perpendicular to each other, is nearly:

- (A)  $i_B = 30^\circ$
- (B)  $i_B = 45^\circ$
- (C)  $i_B = 57^\circ$
- (D)  $i_B = 47^\circ$

**Answer (C)**

**Sol.** From Brewster's law

$$\tan(i_B) = \mu$$

$$i_B = \tan^{-1}(1.5) \approx 57^\circ$$

**Question ID: 102036**

Which of the following is correct for nuclear forces?





- (A) Nuclear forces are much weaker than coulomb force
- (B) The nuclear force between two nucleons falls rapidly to zero as their distance is more than a few femtometre
- (C) Nuclear forces are charge dependent
- (D) Nuclear forces are long range forces

**Answer (B)**

**Sol.** Nuclear forces are strongest, charge independent and very small ranged force.

**Question ID: 102037**

Resistivity ( $\rho$ ) vs temperature graphs of some materials are given below. The graph corresponds to a semiconductor is:



**Answer (C)**

**Sol.** For semiconductor, resistivity decreases with increase in temperature.

**Question ID: 102038**

The length of antenna for transmitting a signal of 30 kHz is:

- (A)  $l = \text{km}$
- (B)  $l = 500 \text{ m}$
- (C)  $l = 2.5 \text{ km}$
- (D)  $l = 10 \text{ km}$

**Answer (C)**

**Sol.**  $l_{\text{min}} = \frac{\lambda}{4} = \frac{c}{4f} = \frac{3 \times 10^8}{4 \times 3 \times 10^4}$

$$= 2500 \text{ m}$$


$$= 2.5 \text{ km}$$

## Question ID: 102039

A conducting rod of mass ' $m$ ' and length ' $l$ ' is changed horizontally with the help of two insulating threads such that the rod is along east-west direction. Now current  $I$  is set up from west to east direction. What should be magnitude and direction of magnetic field. So that tension in the threads becomes zero?

- (A)  $B = \frac{mg}{Il}$ , towards north  
 (B)  $B = \frac{mg}{Il}$ , towards south  
 (C)  $B = \frac{mg}{Il}$ , upwards  
 (D)  $B = \frac{mg}{Il}$ , downwards

## Answer (A)




$$mg = B/L$$

$B = \frac{mg}{Il}$ , towards north to ensure that magnetic force upwards.

## Question ID: 102040

The logic circuit given below will acts like



- (A) NOT gate  
 (B) AND gate  
 (C) OR gate  
 (D) XOR gate

## Answer (C)

Sol.  $Y = \overline{A} \cdot \overline{B} = \overline{A + B} = A + B$

$\Rightarrow$  OR gate

## Question ID: 102041

Read the information given below.

The basic principle of a microwave oven is to generate microwave radiation of appropriate frequency in the working space of the oven where we keep food. This way energy is not wasted in heating up the vessel. In the conventional heating method, the vessel on the burner gets heated first and then the food inside gets heated because of transfer of energy from the vessel. In the microwave oven, on the other hand, energy is directly delivered to water molecules which is shared by the entire food.

The basic principle of a microwave oven is

- (A) Heating effect of current  
 (B) Magnetic effect of current  
 (C) Resonant frequency of water molecule  
 (D) Heat transfer

## Answer (C)

Sol. The basic principle of microwave oven is the resonant frequency of water molecules.

## Question ID: 102042

Read the information given below.

The basic principle of a microwave oven is to generate microwave radiation of appropriate frequency in the working space of the oven where we keep food. This way energy is not wasted in heating up the vessel. In the conventional heating method, the vessel on the burner gets heated first and then the food inside gets heated because of transfer of energy from the vessel. In the microwave oven, on the other hand, energy is directly delivered to water molecules which is shared by the entire food.

Which type of vessel should be used in a microwave ovens?

- (A) Metal container  
 (B) Plastic container  
 (C) Porcelain container  
 (D) Porcelain and metal both

## Answer (C)

Sol. In microwave ovens we prefer porcelain container as they are microwave safe.

**Question ID: 102043**

**Read the information given below.**

The basic principle of a microwave oven is to generate microwave radiation of appropriate frequency in the working space of the oven where we keep food. This way energy is not wasted in heating up the vessel. In the conventional heating method, the vessel on the burner gets heated first and then the food inside gets heated because of transfer of energy from the vessel. In the microwave oven, on the other hand, energy is directly delivered to water molecules which is shared by the entire food.

The frequency range of microwave is

- (A)  $10^7$  Hz to  $10^{19}$  Hz    (B)  $10^9$  Hz to  $10^{12}$  Hz  
(C)  $10^{12}$  Hz to  $10^{15}$  Hz    (D)  $10^{15}$  Hz to  $10^{18}$  Hz

**Answer (B)**

**Sol.** The approximate frequency range of microwave is  $10^9$  Hz to  $10^{12}$  Hz.

**Question ID: 102044**

**Read the information given below.**

The basic principle of a microwave oven is to generate microwave radiation of appropriate frequency in the working space of the oven where we keep food. This way energy is not wasted in heating up the vessel. In the conventional heating method, the vessel on the burner gets heated first and then the food inside gets heated because of transfer of energy from the vessel. In the microwave oven, on the other hand, energy is directly delivered to water molecules which is shared by the entire food.

Microwaves are used in

- (A) Cellular phones    (B) Radar system  
(C) Remote switches    (D) Eye surgery

**Answer (B)**

**Sol.** Microwaves are used in Radar system.

**Question ID: 102045**

**Read the information given below.**

The basic principle of a microwave oven is to generate microwave radiation of appropriate frequency in the working space of the oven where we keep food. This way energy is not wasted in heating up the vessel. In the conventional heating method, the vessel on the burner gets heated first and then the food inside gets heated because of transfer of energy from the vessel. In the microwave oven, on the other hand, energy is directly delivered to water molecules which is shared by the entire food.

Microwaves are detected by:

- (A) Point contact diodes    (B) Photographic film  
(C) Photo cells    (D) Ionisation chamber

**Answer (A)**

**Sol.** Microwaves are detected by point contact diodes.

**Question ID: 102046**

**Read the information given below.**

It is useful to design lenses of derived focal length using surface of suitable radii of curvature. Lens maker formula helps in finding various unknowns and is widely applied.

The radii of curvature of faces of a double concave lens are 20 cm and 30 cm. If refractive index of glass is  $\frac{3}{2}$ , then the focal length of the lens will be:

- (A) 24 cm    (B) -24 cm  
(C) 120 cm    (D) -120 cm

**Answer (B)**

**Sol.** From lens maker's formula

$$\begin{aligned} \frac{1}{f} &= (\mu - 1) \left( \frac{1}{R_1} - \frac{1}{R_2} \right) \\ &= \left( \frac{3}{2} - 1 \right) \left( -\frac{1}{20} - \frac{1}{30} \right) \\ &= \frac{1}{2} \left( -\frac{3-2}{60} \right) = -\frac{1}{24} \\ f &= -24 \text{ cm} \end{aligned}$$

**Question ID: 102047**

**Read the information given below.**

It is useful to design lenses of derived focal length using surface of suitable radii of curvature. Lens maker formula helps in finding various unknowns and is widely applied.

Find the focal length of an equi convex lens whose each face has radius of curvature 12 cm. Given refractive index of glass is  $\frac{3}{2}$ .

- (A) 0    (B) Infinite  $\infty$   
(C) -12 cm    (D) 12 cm

**Answer (D)**

**Sol.** From lens maker's formula

$$\begin{aligned} \frac{1}{f} &= (\mu - 1) \left( \frac{1}{R_1} - \frac{1}{R_2} \right) \\ &= \left( \frac{3}{2} - 1 \right) \left[ \frac{1}{12} - \left( -\frac{1}{12} \right) \right] = \frac{1}{2} \times \frac{2}{12} \\ \Rightarrow f &= 12 \text{ cm} \end{aligned}$$

**Question ID: 102048**

Read the information given below.

It is useful to design lenses of derived focal length using surface of suitable radii of curvature. Lens maker formula helps in finding various unknowns and is widely applied.

A convex lens has 30 cm focal length in air. What is its focal length in a liquid whose refractive index is  $\frac{4}{3}$ ? (Given refractive index glass is  $\frac{3}{2}$ )

- (A) -120 cm                      (B) +120 cm  
(C) 60 cm                      (D) -60 cm

**Answer (B)**

**Sol.**  $\frac{1}{f_{\text{air}}} = (\mu_g - 1) \left( \frac{1}{R_1} - \frac{1}{R_2} \right)$

$$\frac{1}{f_{\text{liquid}}} = \left( \frac{\mu_g}{\mu_l} - 1 \right) \left( \frac{1}{R_1} - \frac{1}{R_2} \right)$$

$$\frac{f_{\text{liq.}}}{f_{\text{air}}} = \frac{1}{1} = 4 \left( \mu_g = \frac{3}{2} \text{ and } \mu_l = \frac{4}{3} \right)$$


$$f_{\text{liq.}} = 4 \times 30 = 120 \text{ cm}$$

**Question ID: 102049**

Read the information given below.

It is useful to design lenses of derived focal length using surface of suitable radii of curvature. Lens maker formula helps in finding various unknowns and is widely applied.

If a convex lens made of a material of refractive index  $n_2$  is immersed in a liquid of refractive index  $n_1$  such that  $n_1 = n_2$ , the incident beam of light will emerge as shown

**Answer (D)**


**Sol.** From lens maker's formula

$$\frac{1}{f} = \left( \frac{\mu_l}{\mu_g} - 1 \right) \left( \frac{1}{R_1} - \frac{1}{R_2} \right)$$

If  $\mu_l = \mu_g$ , then  $\frac{1}{f} = 0$

$$\Rightarrow f = \infty$$

Hence ray diagram will be

**Question ID: 102050**

Read the information given below.

It is useful to design lenses of derived focal length using surface of suitable radii of curvature. Lens maker formula helps in finding various unknowns and is widely applied.

Find the power of a concave lens whose focal length is 40 cm

- (A) -2.5 D                      (B) +2.5 D  
(C)  $+2.5 \times 10^{-2}$  D                      (D)  $-2.5 \times 10^{-2}$  D

**Answer (A)**

**Sol.**  $P = \frac{1}{f(m)} = \frac{1}{(-0.4)}$   
 $= -2.5 \text{ D}$